arXiv:0806.2187v1 [math.AP] 13 Jun 2008

Asymptotic analysis of a boundary-value problem with the nonlinear boundary multiphase interactions in a perforated domain

Taras A. Mel'nyk and Olena A. Sivak

Department of Mathematical Physics, National Taras Shevchenko University of Kyiv,
01033 Kyiv, Ukraine
E-mail: melnyk@imath.kiev.ua, o.a.sivak@gmail.com

Abstract

We consider a boundary-value problem for the second order elliptic differential operator with rapidly oscillating coefficients in a domain Ω_ε that is $\varepsilon-$periodically perforated by small holes. The holes are divided into two $\varepsilon-$periodical sets depending on the boundary interaction at their surfaces. Therefore, two different nonlinear Robin boundary conditions $\sigma_\varepsilon(u_\varepsilon) + \varepsilon\kappa_m(u_\varepsilon) = \varepsilon g_\varepsilon^{(m)},\ m = 1, 2,$ are given on the corresponding boundaries of the small holes. The asymptotic analysis of this problem is made as $\varepsilon \to 0$, namely the convergence theorem both for the solution and for the energy integral is proved without using extension operators, the asymptotic approximations both for the solution and for the energy integral are constructed and the corresponding error estimates are obtained.

1 Introduction and statement of the problem

In recent years, a rich collection of new results on asymptotic analysis of boundary-value problems in perforated domains is appeared (see for example [1]-[10]). The classical method proposed by E. Khruslov [11] and D. Cioranescu and J. Saint Jean Paulin [12] is based on a special bounded extension of solutions in Sobolev spaces. It was established by V. Zhikov [8, 9] that the homogenization results can be obtained without using the extension technique in Sobolev spaces in periodically perforated domains. It should be mentioned the paper [2], where the homogenization results for an elliptic problem with a nonlinear boundary condition in a perforated domain were obtained with the help of a new unfolding method that does not need any extension operators as well.

In this paper we use this simple Zhikov's approach and the scheme of the paper [13], where the full asymptotic analysis (the convergence of the solution and the energy integral, the approximation for the solution and the corresponding asymptotic error estimate in the Sobolev space H^1) was made for an elliptic problem with a nonlinear boundary condition in a thick junction.

Let B be a finite union of smooth disjoint nontangent domains strictly lying in the unit square $\square := \{\xi \in \mathbb{R}^n : \quad 0 < \xi_i < 1, \quad i = \overline{1, n}\}$. In an arbitrary way, we divide B into two sets, $B^{(1)} = \bigcup\limits_{k=1}^{N_1} B_k^{(1)}$ and $B^{(2)} = \bigcup\limits_{k=1}^{N_2} B_k^{(2)}$. Let us introduce the following notations:

$$Q_0 := \square \setminus \overline{B}, \quad \mathcal{B}^{(m)} := \bigcup\nolimits_{z\in\mathbb{Z}^n} \big(z + B^{(m)}\big), \quad \mathcal{B}_\varepsilon^{(m)} := \varepsilon\mathcal{B}^{(m)} = \{x \in \mathbb{R}^n : \ \varepsilon^{-1}x \in \mathcal{B}^{(m)}\},$$

$$m = 1, 2,$$

where ε is a small parameter. Let Ω be a smooth bounded domain in $\mathbb{R}^n$. Define the following perforated domain $\Omega_\varepsilon = \Omega \setminus \overline{\left(\mathcal{B}_\varepsilon^{(1)} \cup \mathcal{B}_\varepsilon^{(2)}\right)}$ and require the domain Ω_ε to be a domain with the Lipschitz boundary. Denote $\Gamma_\varepsilon = \partial\Omega \cap \overline{\Omega_\varepsilon}$ and $\Xi_\varepsilon^{(m)} = \Omega \cap \partial\mathcal{B}_\varepsilon^{(m)}, \quad m = 1, 2, \quad \Xi_\varepsilon = \Xi_\varepsilon^{(1)} \cup \Xi_\varepsilon^{(2)}$. (see Fig. 1).



Figure 1:

Let $a_{ij}(\xi)$, $\xi \in \mathbb{R}^n$, $i, j = \overline{1, n}$, be smooth 1−periodic functions such that

$$\begin{gathered} 1)\ \ \forall i, j = 1, \ldots, n, \quad \forall \xi \in \mathbb{R}^n : \quad a_{ij}(\xi) = a_{ji}(\xi), \\ 2)\ \ \exists\, \varkappa_1 > 0\ \ \exists\, \varkappa_2 > 0 \quad \forall\, \xi \in \mathbb{R}^n \ \ \forall\, \eta \in \mathbb{R}^n : \quad \varkappa_1 |\eta|^2 \le a_{ij}(\xi)\eta_i\eta_j \le \varkappa_2 |\eta|^2. \end{gathered} \tag{1}$$

Remark 1. *Here and in the sequel we adopt the Einstein convention of summation over repeated indexes.*

Let $f_\varepsilon, f_0, g_\varepsilon^{(m)}, g_0^{(m)}$ be given functions such that $f_\varepsilon, f_0 \in L^2(\Omega)$, $g_\varepsilon^{(m)}, g_0^{(m)} \in H_0^1(\Omega)$ and

$$f_\varepsilon \xrightarrow{s} f_0 \quad \text{in} \quad L^2(\Omega), \qquad g_\varepsilon^{(m)} \xrightarrow{w} g_0^{(m)} \quad \text{weakly in} \quad H^1(\Omega), \quad m = 1, 2. \tag{2}$$

The given functions $\kappa_m : \mathbb{R} \to \mathbb{R}$, $m = 1, 2$, are Lipschitz continuous (it is equivalent that $\kappa_m \in W_{loc}^{1,\infty}(\mathbb{R})$) and such that

$$\exists\, c_1 > 0 \ \ \exists\, c_2 > 0 : \ \ c_1 \le \kappa_m' \le c_2 \quad \text{a.e. in } \mathbb{R} \quad (m = 1, 2). \tag{3}$$

In the perforated domain Ω_ε we consider the following nonlinear problem

$$\left\{ \begin{array}{rcll} -\mathcal{L}_\varepsilon(u_\varepsilon) & = & f_\varepsilon & \text{in } \ \Omega_\varepsilon, \\ \sigma_\varepsilon(u_\varepsilon) + \varepsilon\kappa_1(u_\varepsilon) & = & \varepsilon g_\varepsilon^{(1)} & \text{on } \ \Xi_\varepsilon^{(1)}, \\ \sigma_\varepsilon(u_\varepsilon) + \varepsilon\kappa_2(u_\varepsilon) & = & \varepsilon g_\varepsilon^{(2)} & \text{on } \ \Xi_\varepsilon^{(2)}, \\ u_\varepsilon & = & 0 & \text{on } \ \Gamma_\varepsilon, \end{array} \right. \tag{4}$$

where $\mathcal{L}_\varepsilon(u_\varepsilon) \equiv \partial_{x_i}\left(a_{ij}^\varepsilon(x)\partial_{x_j} u_\varepsilon(x)\right)$, $\sigma_\varepsilon(u_\varepsilon) \equiv a_{ij}^\varepsilon(x)\partial_{x_j} u_\varepsilon(x)\,\nu_i$, $a_{ij}^\varepsilon(x) \equiv a_{ij}\left(\frac{x}{\varepsilon}\right)$, $\partial_{x_i} u = \frac{\partial u}{\partial x_i}$, $\left(\nu_1\left(\frac{x}{\varepsilon}\right), \ldots \nu_n\left(\frac{x}{\varepsilon}\right)\right)$ is the outward normal.

Recall that a function u_ε from the Sobolev space $H^1(\Omega_\varepsilon, \Gamma_\varepsilon) = \{u \in H^1(\Omega_\varepsilon) : \ u|_{\Gamma_\varepsilon} = 0\}$ is a weak solution to problem (4) if the following integral identity

$$\int_{\Omega_\varepsilon} a_{ij}^\varepsilon \, \partial_{x_j} u_\varepsilon \, \partial_{x_i} \varphi \, dx + \varepsilon \sum_{m=1}^{2} \int_{\Xi_\varepsilon^{(m)}} \kappa_m(u_\varepsilon) \varphi \, ds_x = \int_{\Omega_\varepsilon} f_\varepsilon \varphi \, dx + \varepsilon \sum_{m=1}^{2} \int_{\Xi_\varepsilon^{(m)}} g_\varepsilon^{(m)} \varphi \, ds_x \qquad (5)$$

holds for any function $\varphi \in H^1(\Omega_\varepsilon, \Gamma_\varepsilon)$.

Our goal is to study the asymptotic behavior of u_ε as $\varepsilon \to 0$. Also it will be understandable further how conduct research in the case of p-multiphase interactions in perforated domains.

2 Auxiliary uniform estimates

Let $H^1_{\text{per}}(Q_0) = \{v \in H^1(Q_0) : \ v - 1\text{-periodic in } \ \xi_1, \ldots, \xi_n\}$. Obviously, we can periodically extend every function v from $H^1_{\text{per}}(Q_0)$ into $H^1_{loc}\big(\mathbb{R}^n \setminus \overline{(\mathcal{B}^{(1)} \cup \mathcal{B}^{(2)})}\big)$; this extension will be denoted again by v. Let $\psi_0^{(m)} \in H^1_{\text{per}}(Q_0)$, $m = 1, 2$, be weak solutions to the corresponding problems

$$\left\{\begin{array}{l} \mathcal{L}_{\xi\xi}\big(\psi_0^{(1)}\big) = q_1 \ \text{ in } \ Q_0, \\ \sigma_\xi\big(\psi_0^{(1)}\big) = 1 \ \text{ on } \ S^{(1)}, \\ \sigma_\xi\big(\psi_0^{(1)}\big) = 0 \ \text{ on } \ S^{(2)}, \\ \langle \psi_0^{(1)} \rangle_{Q_0} = 0, \end{array}\right. \qquad \left\{\begin{array}{l} \mathcal{L}_{\xi\xi}\big(\psi_0^{(2)}\big) = q_2 \ \text{ in } \ Q_0, \\ \sigma_\xi\big(\psi_0^{(2)}\big) = 0 \ \text{ on } \ S^{(1)}, \\ \sigma_\xi\big(\psi_0^{(2)}\big) = 1 \ \text{ on } \ S^{(2)}, \\ \langle \psi_0^{(2)} \rangle_{Q_0} = 0, \end{array}\right. \qquad (6)$$

where $\mathcal{L}_{\xi\xi}(\psi) \equiv \partial_{\xi_i}\left(a_{ij}(\xi) \partial_{\xi_j} \psi(\xi)\right)$, $\sigma_\xi(\psi) \equiv a_{ij} \partial_{\xi_j} \psi(\xi) \nu_i(\xi)$, $(\nu_1, \ldots, \nu_n)$ is the outward normal to S, $S = S^{(1)} \cup S^{(2)}$, $S^{(m)} = \partial B^{(m)}$, $q_m = \frac{|S^{(m)}|}{|Q_0|}$, $|S^{(m)}| = \text{meas}_2 \, S^{(m)}$ $(m = 1, 2)$, $|Q_0| = \text{meas}_3 \, Q_0$, $\langle \psi \rangle_{Q_0} = \int_{Q_0} \psi(\xi) \, d\xi$. The existence and uniqueness of the solutions to problems (6) follows from the lemma.

Lemma 1. *Let* $F_i \in L^2(Q_0), i = \overline{0, n}$, $F_{n+1}^{(m)} \in L^2(S^{(m)})$, $m = 1, 2$. *There exists a solution* $N \in H^1_{\text{per}}(Q_0)$ *to the following problem*

$$\left\{\begin{array}{ll} -\mathcal{L}_{\xi\xi}(N) = F_0 + \partial_{\xi_i} F_i & \textit{in} \quad Q_0, \\ \sigma_\xi(N) = -F_i \nu_i + F_{n+1}^{(1)} & \textit{on} \quad S^{(1)}, \\ \sigma_\xi(N) = -F_i \nu_i + F_{n+1}^{(2)} & \textit{on} \quad S^{(2)}, \end{array}\right. \qquad (7)$$

if and only if

$$\langle F_0 \rangle_{Q_0} + \langle F_{n+1}^{(1)} \rangle_{S^{(1)}} + \langle F_{n+1}^{(2)} \rangle_{S^{(2)}} = 0. \qquad (8)$$

In addition this solution is defined up to an additive constant.

The proof is standard (see for instance [5]). Then the ε-periodic functions $\psi_0^{(m)}\left(\frac{x}{\varepsilon}\right)$, $x \in \Omega_\varepsilon$, $m = 1, 2$, satisfy the following relations

$$\left\{\begin{array}{l} \frac{\partial}{\partial x_i}\big(a_{ij}^\varepsilon(x) \frac{\partial}{\partial x_j}\big(\psi_0^{(1)}\left(\frac{x}{\varepsilon}\right)\big)\big) = \varepsilon^{-2} q_1 \ \text{ in } \Omega_\varepsilon, \\ \sigma_\varepsilon\big(\psi_0^{(1)}\left(\frac{x}{\varepsilon}\right)\big) = \varepsilon^{-1} \ \text{ on } \Xi_\varepsilon^{(1)}, \\ \sigma_\varepsilon\big(\psi_0^{(1)}\left(\frac{x}{\varepsilon}\right)\big) = 0 \ \text{ on } \Xi_\varepsilon^{(2)}, \end{array}\right. \qquad \left\{\begin{array}{l} \frac{\partial}{\partial x_i}\big(a_{ij}^\varepsilon(x) \frac{\partial}{\partial x_j}\big(\psi_0^{(2)}\left(\frac{x}{\varepsilon}\right)\big)\big) = \varepsilon^{-2} q_2 \ \text{ in } \Omega_\varepsilon, \\ \sigma_\varepsilon\big(\psi_0^{(2)}\left(\frac{x}{\varepsilon}\right)\big) = 0 \ \text{ on } \Xi_\varepsilon^{(1)}, \\ \sigma_\varepsilon\big(\psi_0^{(1)}\left(\frac{x}{\varepsilon}\right)\big) = \varepsilon^{-1} \ \text{ on } \Xi_\varepsilon^{(2)}. \end{array}\right.$$

Multiplying with arbitrary function $\varphi \in H^1(\Omega_\varepsilon, \Gamma_\varepsilon)$ the corresponding differential equation, integrating over Ω_ε and taking into account the boundary conditions, we get the following integral identities

$$\varepsilon \int_{\Xi_\varepsilon^{(m)}} \varphi \, ds_x = \varepsilon \int_{\Omega_\varepsilon} a_{ij}^\varepsilon(x)\, \partial_{\xi_j} \psi_0^{(m)}(\xi)|_{\xi=\frac{x}{\varepsilon}} \, \partial_{x_i}\varphi \, dx + q_m \int_{\Omega_\varepsilon} \varphi \, dx \quad m = 1, 2. \tag{9}$$

Due to the regularity properties of solutions to elliptic problems we have

$$\sup_{x\in\Omega_\varepsilon} \left|\nabla_\xi \psi_0^{(m)}(\xi)|_{\xi=\frac{x}{\varepsilon}}\right| = \sup_{\xi\in Q_0} \left|\nabla_\xi \psi_0^{(m)}(\xi)\right| \le C_0 \quad (m = 1, 2). \tag{10}$$

Using Cauchy's inequality with δ $(ab \le \delta a^2 + \frac{b^2}{4\delta}, \ \ a, b, \delta > 0)$ and (10), we deduce from (9) the following estimates (m=1, 2)

$$\varepsilon \int_{\Xi_\varepsilon^{(m)}} \varphi^2 \, ds_x \le C_1 \left(\varepsilon^2 \int_{\Omega_\varepsilon} |\nabla_x \varphi|^2 \, dx + \int_{\Omega_\varepsilon} \varphi^2 \, dx \right), \tag{11}$$

$$\int_{\Omega_\varepsilon} \varphi^2 \, dx \le C_2 \left(\varepsilon^2 \int_{\Omega_\varepsilon} |\nabla_x \varphi|^2 \, dx + \varepsilon \int_{\Xi_\varepsilon^{(m)}} \varphi^2 \, ds_x \right) \qquad \forall\, \varphi \in H^1(\Omega_\varepsilon, \Gamma_\varepsilon), \tag{12}$$

where the constant C_1 and C_2 are independent of ε.

Remark 2. *In what follows all constants $\{C_i\}$ and $\{c_i\}$ in inequalities are independent of the parameter ε.*

It follows from (11) and (2) that

$$\sqrt{\varepsilon} \sum_{m=1}^{2} \|g_\varepsilon^{(m)}\|_{L^2(\Xi_\varepsilon^{(m)})} \le C_3. \tag{13}$$

Also with the help of (11) and (12) it is easy to prove that the usual norm $\|\cdot\|_{H^1(\Omega_\varepsilon)}$ is uniformly equivalent with respect to ε to a new norm

$$\|u\|_\varepsilon := \Big(\int_{\Omega_\varepsilon} |\nabla u|^2 \, dx + \varepsilon \int_{\Xi_\varepsilon} u^2 \, ds_x \Big)^{1/2}$$

in the space $H^1(\Omega_\varepsilon, \Gamma_\varepsilon)$, i.e., there exist constants $C_3 > 0$, $C_4 > 0$ and $\varepsilon_0 > 0$ such that for any $\varepsilon \in (0, \varepsilon_0)$ and $u \in H^1(\Omega_\varepsilon, \Gamma_\varepsilon)$ the following relations hold

$$C_3 \|u\|_{H^1(\Omega_\varepsilon)} \le \|u\|_\varepsilon \le C_4 \|u\|_{H^1(\Omega_\varepsilon)}. \tag{14}$$

2.1 Existence and uniqueness of the solution to problem (4)

Associated with (4), we consider the energy functional

$$I_\varepsilon[u] := \tfrac{1}{2} \int\limits_{\Omega_\varepsilon} a_{ij}^\varepsilon(x)\, \partial_{x_i} u \, \partial_{x_j} u \, dx + \varepsilon \sum_{m=1}^{2} \Big(\int\limits_{\Xi_\varepsilon^{(m)}} K^{(m)}(u) \, ds_x - \int\limits_{\Xi_\varepsilon^{(m)}} g_\varepsilon^{(m)} u \, ds_x \Big) - \int\limits_{\Omega_\varepsilon} f_\varepsilon u \, dx \tag{15}$$

on $H^1(\Omega_\varepsilon, \Gamma_\varepsilon)$, where

$$K^{(m)}(z) = \int_0^z \kappa_m(t) \, dt \quad \forall\, z \in \mathbb{R}, \quad m = 1, 2. \tag{16}$$

It is easy to prove that if u_ε is a minimizer of I_ε at a fixed value of ε, then u_ε is a weak solution to problem (4).

Theorem 1. *At each fixed value of ε problem (4) has exactly one solution $u_\varepsilon \in H^1(\Omega_\varepsilon, \Gamma_\varepsilon)$ for which the following estimate*

$$\|u_\varepsilon\|_{H^1(\Omega_\varepsilon)} \le C_1\Big(1 + \|f_\varepsilon\|_{L^2(\Omega_\varepsilon)} + \sqrt{\varepsilon}\sum\nolimits_{m=1}^{2} \|g_\varepsilon^{(m)}\|_{L^2(\Xi_\varepsilon^{(m)})}\Big) \le C_2 \tag{17}$$

holds, where the constants C_1 and C_2 are independent of ε, $f_\varepsilon, g_\varepsilon^{(m)}$ and u_ε.

Proof. Integrating inequalities in (3), we obtain

$$c_1 t^2 + \kappa_m(0)\, t \le \kappa_m(t)\, t \le c_2 t^2 + \kappa_m(0)\, t \quad \forall\, t \in \mathbb{R}, \tag{18}$$

whence it follows that

$$\frac{c_1}{2}\, z^2 + \kappa_m(0) z \le K^{(m)}(z) \le \frac{c_2}{2}\, z^2 + \kappa_m(0) z \quad \forall\, z \in \mathbb{R} \quad m = 1, 2. \tag{19}$$

Using (14), (18), (19), (2) and the same arguments as in Theorem 1 ([13]), we can prove the coercitivity condition on I, i.e., the following inequality

$$I_\varepsilon[u] \ge C_1 \|u\|^2_{H^1(\Omega_\varepsilon)} - C_2 \tag{20}$$

holds for any function $u \in H^1(\Omega_\varepsilon, \Gamma_\varepsilon)$.

With the help of (9) we can re-write the energy functional as

$$\begin{aligned} I_\varepsilon[u] = \frac{1}{2}\int_{\Omega_\varepsilon} a_{ij}^\varepsilon(x)\, \partial_{x_i} u\, \partial_{x_j} u\, dx + \sum_{m=1}^{2}\Big(\varepsilon \int_{\Omega_\varepsilon} a_{ij}^\varepsilon\, \partial_{\xi_j}\big(\psi_0(\xi)\big)\big|_{\xi=\frac{x}{\varepsilon}}\, \kappa_m(u)\, \partial_{x_i} u\, dx + \\ + q_m \int\limits_{\Omega_\varepsilon} K^{(m)}(u)\, dx - \varepsilon \int\limits_{\Omega_\varepsilon} a_{ij}^\varepsilon \partial_{\xi_j}\big(\psi_0(\xi)\big)\big|_{\xi=\frac{x}{\varepsilon}}\big(u \partial_{x_i} g_\varepsilon^{(m)} + g_\varepsilon^{(m)} \partial_{x_i} u\big)\, dx - \\ - q_m \int\limits_{\Omega_\varepsilon} g_\varepsilon^{(m)} u\, dx\Big) - \int\limits_{\Omega_\varepsilon} f_\varepsilon u\, dx. \end{aligned}$$

Consider the function

$$\begin{aligned} L(p, t, x) = \frac{1}{2} a_{ij}^\varepsilon p_i p_j + \sum_{m=1}^{2}\Big(\varepsilon a_{ij}^\varepsilon\, \partial_{\xi_j}\big(\psi_0(\xi)\big)\big|_{\xi=\frac{x}{\varepsilon}}\, \kappa_m(t)\, p_i + q_m K^{(m)}(t) - \\ - \varepsilon a_{ij}^\varepsilon \partial_{\xi_j}\big(\psi_0(\xi)\big)\big|_{\xi=\frac{x}{\varepsilon}}\big(t \partial_{x_i} g_\varepsilon^{(m)} + g_\varepsilon^{(m)} p_i\big) - q_m t g_\varepsilon^{(m)}\Big) - f_\varepsilon t. \end{aligned}$$

Since

$$\partial^2_{p_i p_j} L(p, t, x)\, \eta_i \eta_j = 2^{-1} a_{ij}^\varepsilon(x) \eta_i \eta_j \ge \varkappa_1 |\eta|^2 \quad \forall\, p, \eta \in \mathbb{R}^n, \quad x \in \Omega_\varepsilon,$$

the function L is uniformly convex in p for each $x \in \Omega_\varepsilon$. This means that $I[\cdot]$ is weakly lower semicontinuous on $H^1(\Omega_\varepsilon, \Gamma_\varepsilon)$ and there exists at least one minimizer (see [14, Chapter 8.2]).

Thanks to (3) it is easy to prove the uniqueness of this minimizer (see Theorem 1 ([13])).

Finally, let us deduce the uniform estimate (17). Denote by u_ε the solution to problem (4). Setting $\varphi = u_\varepsilon$ in (5) and taking into account (1) and the left inequality in (18), we get

$$\varkappa_1 \int_{\Omega_\varepsilon} |\nabla u_\varepsilon|^2\, dx + \varepsilon\, c_1 \int_{\Xi_\varepsilon} u_\varepsilon^2\, ds_x + \varepsilon \kappa_m(0) \sum_{m=1}^{2} \int_{\Xi_\varepsilon^{(m)}} u_\varepsilon\, ds_x \le \int_{\Omega_\varepsilon} f_\varepsilon u_\varepsilon\, dx + \varepsilon \sum_{m=1}^{2} \int_{\Xi_\varepsilon^{(m)}} g_\varepsilon^{(m)} u_\varepsilon\, ds_x$$

from which

$$c_2\Big(\int_{\Omega_\varepsilon}|\nabla u_\varepsilon|^2\,dx+\varepsilon\int_{\Xi_\varepsilon}u_\varepsilon^2\,d\sigma_x\Big)\le c_3\sqrt{\varepsilon}\|u_\varepsilon\|_{L^2(\Xi_\varepsilon)}+\|f_\varepsilon\|_{L^2(\Omega_\varepsilon)}\|u_\varepsilon\|_{L^2(\Omega_\varepsilon)}+$$
$$+\,\varepsilon\sum_{m=1}^{2}\|g_\varepsilon^{(m)}\|_{L^2(\Xi_\varepsilon^{(m)})}\|u_\varepsilon\|_{L^2(\Xi_\varepsilon^{(m)})}.$$

Using (14) and (11), we derive the first part of the estimate (17) from the last inequality, and then the second one on the basis of (2) and (13). □

3 Convergence theorem

In the sequel, $\widetilde{y}$ denotes the zero-extension of a function y defined on Ω_ε into the domain Ω. Also we introduce the following characteristic function

$$\chi_{Q_0}(\xi)=\begin{cases}1, & x\in Q_0,\\ 0, & x\in\square\setminus Q_0.\end{cases} \tag{21}$$

It is known that $\chi^\varepsilon_{Q_0}(x):=\chi_{Q_0}(\frac{x}{\varepsilon})\xrightarrow{w}|Q_0|$ weakly in $L^2(\Omega)$ as $\varepsilon\to0$.

Lemma 2. *Let $\{v_\varepsilon\}_{\varepsilon>0}$ be a sequence in $H^1(\Omega_\varepsilon,\Gamma_\varepsilon)$ uniformly bounded in ε in $H^1(\Omega_\varepsilon,\Gamma_\varepsilon)$ and such that*

$$\widetilde{\kappa_m(v_\varepsilon)}\to\zeta\quad\text{weakly in }\ L^2(\Omega)\quad\text{as}\quad\varepsilon\to0\qquad(m=1,2).$$

Then for any function $\varphi\in H^1(\Omega_\varepsilon,\Gamma_\varepsilon)$

$$\varepsilon\int_{\Xi_\varepsilon^{(m)}}\kappa_m(v_\varepsilon)\,\varphi\,ds_x\ \to\ q_m\int_\Omega\zeta(x)\,\varphi(x)\,dx\quad\text{as}\quad\varepsilon\to0\qquad(m=1,2). \tag{22}$$

Proof. By virtue of (9) we have

$$\varepsilon\int_{\Xi_\varepsilon^{(m)}}\kappa(v_\varepsilon)\,\varphi\,ds_x=\varepsilon\int_{\Omega_\varepsilon}a_{ij}^\varepsilon(x)\,\partial_{\xi_j}\psi_0^{(m)}(\xi)|_{\xi=\frac{x}{\varepsilon}}\big(\kappa'(v_\varepsilon)\,\partial_{x_i}v_\varepsilon\,\varphi+\kappa(v_\varepsilon)\,\partial_{x_i}\varphi\big)\,dx+$$
$$+\,q_m\int_\Omega\widetilde{\kappa(v_\varepsilon)}\,\varphi\;dx,\quad m=1,2.$$

Thanks to the Lemma's condition, (3) and (10), the first summand vanishes and the second one tends to $q_m\int_\Omega\zeta(x)\,\varphi\,dx$ as $\varepsilon\to0\ \ m=1,2.$ □

Remark 3. *From Lemma 2 it follows that for any sequence $\{v_\varepsilon\}_{\varepsilon>0}\in H^1(\Omega_\varepsilon,\Gamma_\varepsilon)$, which is uniformly bounded with respect to ε, there exists a subsequence $\{\varepsilon'\}\subset\{\varepsilon\}$ (again denoted by $\{\varepsilon\}$) and a function $\zeta\in L^2(\Omega)$ such that the convergences (22) hold.*

Using (2), we can prove similarly as in Lemma 2 that for any function $\varphi\in H^1(\Omega_\varepsilon,\Gamma_\varepsilon)$

$$\varepsilon\int_{\Xi_\varepsilon^{(m)}}g_\varepsilon^{(m)}\varphi\,ds_x\ \to\ |S^{(m)}|\int_\Omega g_0^{(m)}(x)\,\varphi(x)\,dx\quad\text{as}\quad\varepsilon\to0\qquad(m=1,2). \tag{23}$$

Consider $1-$periodic solutions $T_l,\ l=1,\dots,n$, to the following problems

$$\left\{ \begin{array}{l} \mathcal{L}_{\xi\xi}\big(T_l\big) = -\partial_{\xi_i} a_{il} \ \text{ in } \ Q_0, \\ \sigma_\xi\big(T_l\big) = -a_{il}\,\nu_i \ \text{ on } \ S, \quad \langle T_l\rangle_{Q_0} = 0. \end{array} \right. \tag{24}$$

From Lemma 1 it follows the existence and uniqueness of the solutions to these problems.

With the help of $T_l,\ l=1,\dots,n$, we define the coefficients of the homogenized matrix $\{\widehat{a}_{ij}\}$ by the formula

$$\widehat{a}_{ij} = \langle a_{ij} + a_{ik}\partial_{\xi_k} T_j\rangle_{Q_0}, \quad i,j\in\{1,2,\dots,n\}. \tag{25}$$

It is easy to see that

$$\widehat{a}_{ij} = \langle a_{kl}\,\partial_{\xi_k}(\xi_i + T_i)\,\partial_{\xi_l}(\xi_j + T_j)\rangle_{Q_0} \tag{26}$$

i.e., the matrix $\{\widehat{a}_{ij}\}$ is symmetric and it is well known that it is elliptic (see for instance [5]).

Theorem 2. *For the solution u_ε to problem (4) there exists the following convergences*

$$\left.\begin{array}{rcll} \widetilde{u_\varepsilon} & \xrightarrow{w} & |Q_0|\,v_0 & \textit{weakly in } \ L^2(\Omega), \\ \widetilde{a^\varepsilon_{ij}\,\partial_{x_j} u_\varepsilon} & \xrightarrow{w} & \widehat{a}_{ij}\partial_{x_j} v_0 & \textit{weakly in } \ L^2(\Omega), \quad i=1,\dots,n, \end{array}\right\} \ \textit{as} \ \ \varepsilon\to 0, \tag{27}$$

where v_0 is a unique weak solution to the following problem

$$\left\{ \begin{array}{rcll} -\widehat{a}_{ij}\,\partial^2_{x_ix_j} v_0(x) + \sum\limits_{m=1}^{2} |S^{(m)}|\,\kappa_m(v_0(x)) & = & \sum\limits_{m=1}^{2} |S^{(m)}|\,g_0^{(m)}(x) + |Q_0|\,f_0(x), & x\in\Omega, \\ v_0(x) & = & 0, & x\in\partial\Omega, \end{array} \right. \tag{28}$$

which is called homogenized problem for (4).

Furthermore, the following energy convergence holds as $\varepsilon\to 0$:

$$E_\varepsilon(u_\varepsilon) := \int_{\Omega_\varepsilon} a^\varepsilon_{ij}(x)\,\partial_{x_j} u_\varepsilon\,\partial_{x_i} u_\varepsilon\,dx + \varepsilon\sum_{m=1}^{2}\int_{\Xi^{(m)}_\varepsilon} \kappa_m(u_\varepsilon)\,u_\varepsilon\,ds_x \longrightarrow$$

$$\int_\Omega \widehat{a}_{ij}\,\partial_{x_j} v_0\,\partial_{x_i} v_0\,dx \ + \sum_{m=1}^{2}|S^{(m)}|\int_\Omega \kappa_m(v_0)\,v_0\,dx =: E_0(v_0). \tag{29}$$

Proof. **1.** It follows from (17) and (3) that the values

$$\|\widetilde{u_\varepsilon}\|_{L^2(\Omega)}, \quad \|\widetilde{a^\varepsilon_{ij}\,\partial_{x_j} u_\varepsilon}\|_{L^2(\Omega)}, \ \ i=1,\dots,n, \quad \|\widetilde{\kappa_m(u_\varepsilon)}\|_{L^2(\Omega)}, \ \ m=1,2,$$

are uniformly bounded with respect to ε. Hence there exists a subsequence $\{\varepsilon'\}\subset\{\varepsilon\}$, again denoted by $\{\varepsilon\}$, such that

$$\left.\begin{array}{rcll} \widetilde{u_\varepsilon} & \xrightarrow{w} & |Q_0|\,v_0 & \text{weakly in } \ L^2(\Omega), \\ \widetilde{a^\varepsilon_{ij}\,\partial_{x_j} u_\varepsilon} & \xrightarrow{w} & \gamma_i & \text{weakly in } \ L^2(\Omega), \quad i=1,\dots,n, \\ \widetilde{\kappa_m(u_\varepsilon)} & \xrightarrow{w} & \zeta_m & \text{weakly in } \ L^2(\Omega), \quad m=1,2, \end{array}\right\} \ \text{as} \ \ \varepsilon\to 0, \tag{30}$$

where $v_0,\ \gamma_i,\ i=1,\dots,n,\ \zeta_m,\ m=1,2$, are some functions which will be determined in what follows.

2. Obviously the ε-periodic functions $T_l\left(\frac{\cdot}{\varepsilon}\right),\ l = 1, \ldots, n$, defined in (24) satisfy the following relations

$$\begin{cases} \partial_{x_i}\left(a_{ij}(\xi)\partial_{\xi_j}T_l(\xi)|_{\xi=\frac{x}{\varepsilon}}\right) + \partial_{x_i}a^{\varepsilon}_{il}(x) = 0 \quad \forall\, x \in \Omega_\varepsilon, \\ \big(a_{ij}(\xi)\,\partial_{\xi_j}T_l(\xi)\,\nu_i(\xi) + a_{il}(\xi)\nu_i(\xi)\big)\Big|_{\xi=\frac{x}{\varepsilon}} = 0 \quad \forall\, x \in \Xi_\varepsilon. \end{cases}$$

Multiplying the first relation by $u_\varepsilon\,\phi$, where ϕ is arbitrary function from $C_0^\infty(\Omega)$, and integrating over Ω_ε, we obtain

$$\int_{\Omega_\varepsilon}\big(a_{ij}(\xi)\,\partial_{\xi_j}T_l(\xi) + a_{il}(\xi)\big)|_{\xi=\frac{x}{\varepsilon}}\,\big(u_\varepsilon\,\partial_{x_i}\phi + \phi\,\partial_{x_i}u_\varepsilon\big)\,dx = 0, \qquad l = \overline{1,n}. \tag{31}$$

Put the following test-function $\varphi(x) = \varepsilon T_l(\frac{x}{\varepsilon})\phi(x),\ x \in \Omega_\varepsilon$, into the integral identity (5). The result is as follows

$$\int_{\Omega_\varepsilon} a^{\varepsilon}_{ij}(x)\partial_{x_j}u_\varepsilon\partial_{\xi_i}T_l(\xi)|_{\xi=\frac{x}{\varepsilon}}\,\phi(x)\,dx + \varepsilon\int_{\Omega_\varepsilon} a^{\varepsilon}_{ij}(x)\partial_{x_j}u_\varepsilon\,T_l(\frac{x}{\varepsilon})\,\partial_{x_i}\phi(x)\,dx +$$
$$+\,\varepsilon^2\sum_{m=1}^{2}\int_{\Xi^{(m)}_\varepsilon}\kappa_m(u_\varepsilon)\,T_l\,\phi\,ds_x = \varepsilon\int_{\Omega_\varepsilon} f_\varepsilon\,T_l\,\phi\,dx + \varepsilon^2\sum_{m=1}^{2}\int_{\Xi^{(m)}_\varepsilon} g^{(m)}_\varepsilon\,T_l\,\phi\,ds_x. \tag{32}$$

Using (2), (3) and the identities (9), it follows from (32) that

$$\int_{\Omega_\varepsilon} a^{\varepsilon}_{ij}(x)\partial_{x_j}u_\varepsilon\partial_{\xi_i}T_l(\xi)|_{\xi=\frac{x}{\varepsilon}}\,\phi(x)\,dx = \mathcal{O}(\varepsilon) \quad \text{as} \quad \varepsilon \to 0, \quad l = \overline{1,n}. \tag{33}$$

Subtracting (32) from (31), we get

$$\int_{\Omega}\big(a_{ij}(\xi)\,\partial_{\xi_j}T_l(\xi) + a_{il}(\xi)\big)|_{\xi=\frac{x}{\varepsilon}}\,\widetilde{u_\varepsilon}\,\partial_{x_i}\phi\,dx + \int_{\Omega}\widetilde{a^{\varepsilon}_{il}\,\partial_{x_i}u_\varepsilon}\,\phi\,dx = \mathcal{O}(\varepsilon), \qquad l = \overline{1,n}. \tag{34}$$

In (34) we regard that the functions $a_{ij}\,\partial_{\xi_j}T_l + a_{il},\ l = 1, \ldots, n$, are equal to zero on B.

Let us find the limit of the first summand in the left-hand side of (34). At first we note that the limit function v_0 in (30) belongs to $H^1_0(\Omega)$ because of the conectedness of the domain $\mathbb{R}^n \setminus \overline{\big(\mathcal{B}^{(1)} \cup \mathcal{B}^{(2)}\big)}$ (see [8]-[10]). Since $\big(a_{ij}(\xi)\,\partial_{\xi_j}T_l(\xi) + a_{il}(\xi)\big)\nu_i(\xi) = 0$ at $\xi \in S$ and the vector-functions

$$\mathbf{F}_l = \big(a_{1j}(\xi)\,\partial_{\xi_j}T_l(\xi) + a_{1l}(\xi)\,, \ldots,\ a_{nj}(\xi)\,\partial_{\xi_j}T_l(\xi) + a_{nl}(\xi)\big), \quad l = 1, \ldots, n, \tag{35}$$

are solenoidal in Q_0 (see (24)), their zero-extensions into $\square \setminus Q_0$ are also solenoidal in weak sense, i.e.,

$$\int_{Q_0}\mathbf{F}_l(\xi)\cdot\nabla\psi(\xi)\,d\xi = \int_{\square}\mathbf{F}_l(\xi)\cdot\nabla\psi(\xi)\,d\xi = 0 \quad \forall\psi \in C^\infty_{\text{per}}(\square), \quad l = 1, \ldots, n.$$

Then using results by V.V. Zhikov (see [8, Th. 2.1]), we get that

$$\lim_{\varepsilon\to 0}\int_{\Omega}\big(a_{ij}(\xi)\,\partial_{\xi_j}T_l(\xi) + a_{il}(\xi)\big)|_{\xi=\frac{x}{\varepsilon}}\,\widetilde{u_\varepsilon}\,\partial_{x_i}\phi\,dx = \int_{\Omega}\widehat{a}_{il}\,v_0\,\partial_{x_i}\phi\,dx.$$

As a results, it follows from (34) in the limit passage as $\varepsilon \to 0$ that

$$\int_\Omega \widehat{a}_{il}\, v_0\, \partial_{x_i}\phi\, dx + \int_\Omega \gamma_l\, \phi\, dx = 0 \quad \forall \phi \in C_0^\infty(\Omega), \quad (l = 1, \ldots, n),$$

i.e.,

$$\gamma_l(x) = \widehat{a}_{il}\, \partial_{x_i} v_0(x) \quad \text{for a.e. } x \in \Omega \quad (l = 1, \ldots, n). \tag{36}$$

4. Using the extension by zero and the identities (9), we rewrite the integral identity (5) in the following way

$$\begin{gathered}\int_\Omega \widetilde{a_{ij}^\varepsilon\, \partial_{x_j} u_\varepsilon}\, \partial_{x_i}\varphi\, dx + \\ + \sum_{m=1}^{2} \Bigg(\underbrace{\varepsilon \int_{\Omega_\varepsilon} a_{ij}^\varepsilon(x)\, \partial_{\xi_j}\psi_0^{(m)}(\xi)|_{\xi=\frac{x}{\varepsilon}} \left(\kappa_m'(u_\varepsilon)\, \partial_{x_i} u_\varepsilon\, \varphi + \kappa_m(u_\varepsilon)\, \partial_{x_i}\varphi\right) dx} + q_m \int_\Omega \widetilde{\kappa_m(u_\varepsilon)}\, \varphi\, dx \Bigg) = \\ = \int_\Omega \chi_{Q_0}^\varepsilon\, f_\varepsilon \varphi\, dx + \sum_{m=1}^{2} \Bigg(\underbrace{\varepsilon \int_{\Omega_\varepsilon} a_{ij}^\varepsilon(x)\, \partial_{\xi_j}\psi_0^{(m)}(\xi)|_{\xi=\frac{x}{\varepsilon}} \left(\partial_{x_i} g_\varepsilon^{(m)}\, \varphi + g_\varepsilon^{(m)}\, \partial_{x_i}\varphi\right) dx} + \\ + q_m \int_\Omega \chi_{Q_0}^\varepsilon\, g_\varepsilon^{(m)}\, \varphi\, dx \Bigg) \quad \forall\, \varphi \in C_0^\infty(\Omega).\end{gathered} \tag{37}$$

It is easy to see that the pointed summands in (37) vanish as $\varepsilon \to 0$; the first one due to (3), (10) and (17), the second one due to (10) and (2).

Taking into account (30), (36) and (2), we pass to the limit in (37) as $\varepsilon \to 0$. As a result we get the identity

$$\int_\Omega \widehat{a}_{ij}\, \partial_{x_j} v_0\, \partial_{x_i}\varphi\, dx + \sum_{m=1}^{2} q_m \int_\Omega \zeta_m\, \varphi\, dx = |Q_0| \int_\Omega f_0 \varphi\, dx + \sum_{m=1}^{2} |S^{(m)}| \int_\Omega g_0^{(m)}\, \varphi\, dx \tag{38}$$

for any function $\varphi \in C_0^\infty(\Omega)$. Since the space $C_0^\infty(\Omega)$ is dense in $H_0^1(\Omega)$, identity (38) is valid for any function $\varphi \in H_0^1(\Omega)$.

5. With the help of (2), (5) and (38) we can find that

$$\begin{gathered}\lim_{\varepsilon\to 0} E_\varepsilon(u_\varepsilon) = \lim_{\varepsilon\to 0} \left(\int_{\Omega_\varepsilon} f_\varepsilon\, u_\varepsilon\, dx + \sum_{m=1}^{2} \int_{\Xi_\varepsilon^{(m)}} g_\varepsilon^{(m)}\, u_\varepsilon\, ds_x \right) = \lim_{\varepsilon\to 0} \Bigg(\int_\Omega f_\varepsilon\, \widetilde{u_\varepsilon}\, dx + \\ + \sum_{m=1}^{2} \left(\varepsilon \int_{\Omega_\varepsilon} a_{ij}^\varepsilon(x)\, \partial_{\xi_j}\psi_0^{(m)}|_{\xi=\frac{x}{\varepsilon}} \left(\partial_{x_i} g_\varepsilon^{(m)}\, u_\varepsilon + g_\varepsilon^{(m)}\, \partial_{x_i} u_\varepsilon\right) dx + q_m \int_\Omega g_\varepsilon^{(m)}\, \widetilde{u_\varepsilon}\, dx \right) \Bigg) = \\ = |Q_0| \int_\Omega f_0 v_0\, dx + \sum_{m=1}^{2} |S^{(m)}| \int_\Omega g_0^{(m)} v_0\, dx = \int_\Omega \widehat{a}_{ij} \partial_{x_j} v_0 \partial_{x_i} v_0\, dx + \sum_{m=1}^{2} q_m \int_\Omega \zeta_m(x) v_0\, dx.\end{gathered} \tag{39}$$

6. Now it remains to determine the last summand in (39). For this we will use the method of Browder and Minty, a remarkable technique which somehow applies to the corresponding inequality of monotonicity to justify passing to a weak limit within a nonlinearity.

Thanks to (1) and (3), the inequality of monotonicity in our case reads as follows

$$\int_{\Omega_\varepsilon} a^\varepsilon_{ij}\,\partial_{x_j}\big(u_\varepsilon-\varphi-\varepsilon T_p\,\partial_{x_p}\varphi\big)\,\partial_{x_i}\big(u_\varepsilon-\varphi-\varepsilon T_q\,\partial_{x_q}\varphi\big)\,dx+$$
$$+\,\varepsilon\sum_{m=1}^{2}\int_{\Xi^{(m)}_\varepsilon}\big(\kappa_m(u_\varepsilon)-\kappa_m(\varphi)\big)\big(u_\varepsilon-\varphi\big)\,ds_x\ge 0\qquad\forall\,\varphi\in C^\infty_0(\Omega),\tag{40}$$

which is equivalent to

$$\int_{\Omega_\varepsilon} a^\varepsilon_{ij}\,\partial_{x_j}u_\varepsilon\,\partial_{x_i}u_\varepsilon\,dx+\varepsilon\sum_{m=1}^{2}\int_{\Xi^{(m)}_\varepsilon}\kappa_m(u_\varepsilon)\,u_\varepsilon\,ds_x+$$
$$+\int_{\Omega_\varepsilon} a^\varepsilon_{ij}\,\big(\partial_{x_j}\varphi+\partial_{\xi_j}T_p\,\partial_{x_p}\varphi\big)\,\big(\partial_{x_i}\varphi+\partial_{\xi_i}T_q\,\partial_{x_q}\varphi\big)\,dx-2\int_{\Omega}\widetilde{a^\varepsilon_{ij}\,\partial_{x_j}u_\varepsilon}\,\partial_{x_i}\varphi\,dx-$$
$$-\,2\int_{\Omega_\varepsilon} a^\varepsilon_{ij}\,\partial_{x_j}u_\varepsilon\,\partial_{\xi_i}T_q\,\partial_{x_q}\varphi\,dx-2\varepsilon\int_{\Omega_\varepsilon} a^\varepsilon_{ij}\,\big(\partial_{x_j}u_\varepsilon-\partial_{x_j}\varphi-\partial_{\xi_j}T_p\,\partial_{x_p}\varphi\big)\,T_q\,\partial^2_{x_ix_q}\varphi\,dx+$$
$$+\,\varepsilon^2\int_{\Omega_\varepsilon} a^\varepsilon_{ij}\,T_p\,T_q\,\partial^2_{x_jx_p}\varphi\,\partial^2_{x_ix_q}\varphi\,dx-$$
$$-\,\varepsilon\sum_{m=1}^{2}\int_{\Xi^{(m)}_\varepsilon}\big(\kappa_m(\varphi)\,u_\varepsilon+\kappa_m(u_\varepsilon)\,\varphi-\kappa_m(\varphi)\,\varphi\big)\,ds_x\ge 0\quad\forall\,\varphi\in C^\infty_0(\Omega).\tag{41}$$

The limit of the first line in (41) is equal to the right-hand side in (39). The first integral in the second line can be re-written in the form

$$\int_{\Omega_\varepsilon}\Big(a_{ij}(\xi)\,\partial_{\xi_j}\big(\xi_p+T_p\big)\partial_{\xi_i}\big(\xi_q+T_q\big)\Big)|_{\xi=\frac{x}{\varepsilon}}\,\partial_{x_p}\varphi\,\partial_{x_q}\varphi\,dx.\tag{42}$$

It follows from [8] that its limit equals $\int_\Omega \widehat{a}_{pq}\,\partial_{x_p}\varphi\,\partial_{x_q}\varphi\,dx$. Due to (33) the integral in third line vanishes. Obviously, the limits of summands in the fourth line are equal to zero. The limits of the integrals in the last line can be found with the help of Lemma 2. As a results we have

$$\int_{\Omega}\widehat{a}_{ij}\,\partial_{x_j}(v_0-\varphi)\,\partial_{x_i}(v_0-\varphi)\,dx+\sum_{m=1}^{2}q_m\int_{\Omega}\big(\zeta_m-|Q_0|\kappa_m(\varphi)\big)\,\big(v_0-\varphi\big)\,dx\ge 0.\tag{43}$$

Evidently, this inequality holds for any function $\varphi\in H^1_0(\Omega)$.

Fix any $\psi\in C^\infty_0(\Omega)$ and set $\varphi:=v_0-\lambda\psi$ ($\lambda>0$) in (43). We get then

$$\lambda\int_{\Omega}\widehat{a}_{ij}\,\partial_{x_j}\psi\,\partial_{x_i}\psi\,dx+\sum_{m=1}^{2}q_m\int_{\Omega}\big(\zeta_m-|Q_0|\kappa_m(v_0-\lambda\psi)\big)\,\psi\,dx\ge 0\quad\forall\,\psi\in C^\infty_0(\Omega).$$

In the limit (as $\lambda\to 0$) we obtain

$$\int_{\Omega}\sum_{m=1}^{2}q_m\big(\zeta_m-|Q_0|\kappa_m(v_0)\big)\,\psi\,dx\ge 0.$$

Replacing ψ by $-\psi$, we deduce that in fact quality holds above. Thus

$$\sum_{m=1}^{2} q_m \zeta_m(x) = \sum_{m=1}^{2} |S^{(m)}|\, \kappa_m(v_0(x)) \quad \text{for a.e. } x \in \Omega. \tag{44}$$

7. Returning to (38), we see that the function v_0 satisfies the following integral identity

$$\int_\Omega \widehat{a}_{ij}\, \partial_{x_j} v_0\, \partial_{x_i}\varphi\, dx + \sum_{m=1}^{2} |S^{(m)}| \int_\Omega \kappa_m(v_0)\, \varphi\, dx = |Q_0| \int_\Omega f_0 \varphi\, dx + \sum_{m=1}^{2} |S^{(m)}| \int_\Omega g_0^{(m)}\, \varphi\, dx \tag{45}$$

for any function $\varphi \in H_0^1(\Omega)$. Hence v_0 is a weak solution to the limit problem (28). Thanks to (3) this solution is unique.

Due to the uniqueness of the solution to problem (28), the above argumentations hold for any subsequence of $\{\varepsilon\}$ chosen at the beginning of the proof. By replacing (44) in (39), one obtains the convergence of energies (29). □

4 Asymptotic approximation to the solution and the energy integral

We take the following approximation

$$\overline{u}_\varepsilon := v_0(x) + \varepsilon T_k\left(\tfrac{x}{\varepsilon}\right) \partial_{x_k} v_0(x) \tag{46}$$

to the solution u_ε. Substituting the difference $u_\varepsilon - \overline{u}_\varepsilon$, we find the residuals both in the differential equation and boundary conditions. Straightforward calculation show that

$$\begin{aligned} -\mathcal{L}_\varepsilon\left(u_\varepsilon - \overline{u}_\varepsilon\right) &= f_\varepsilon(x) - f_0(x) - \sum_{m=1}^{2} q_m \left(g_0^{(m)}(x) - \kappa_m(v_0(x))\right) + \\ &+ \left(a_{ij}(\xi) + a_{ik}(\xi)\partial_{\xi_k} T_j(\xi) - \tfrac{1}{|Q_0|}\, \widehat{a}_{ij}\right)\Big|_{\xi=\frac{x}{\varepsilon}}\ \partial^2_{x_i x_j} v_0 + \varepsilon \partial_{x_i}\left(F_i^\varepsilon(x)\right), \quad x \in \Omega_\varepsilon; \end{aligned} \tag{47}$$

$$\sigma_\varepsilon\left(u_\varepsilon - \overline{u}_\varepsilon\right) = -\varepsilon \kappa_m(u_\varepsilon) + \varepsilon g_\varepsilon^{(m)}(x) - F_i^\varepsilon(x)\, \nu_i, \quad x \in \Xi_\varepsilon^{(m)} \quad (m = 1, 2), \tag{48}$$

where $F_i^\varepsilon(x) = a_{ij}(\frac{x}{\varepsilon}) T_k(\frac{x}{\varepsilon})\, \partial^2_{x_j x_k} v_0(x)$, $i = 1, \ldots, n$, and

$$(u_\varepsilon - \overline{u}_\varepsilon)|_{\Gamma_\varepsilon} = -\varepsilon T_k\left(\tfrac{x}{\varepsilon}\right) \partial_{x_k} v_0(x). \tag{49}$$

Let φ_ε be a smooth function in $\overline{\Omega}$ such that $0 \le \varphi_\varepsilon \le 1$, $\varphi_\varepsilon(x) = 1$ if $dist(x, \partial\Omega) \le \varepsilon$, and $\varphi_\varepsilon(x) = 0$ if $dist(x, \partial\Omega) \ge 2\varepsilon$. Obviously,

$$|\nabla_x \varphi_\varepsilon| \le c\, \varepsilon^{-1} \quad \text{in } \overline{\Omega}. \tag{50}$$

With the help of φ_ε we define the following functions

$$\psi_\varepsilon(x) = -\varepsilon \varphi_\varepsilon(x)\, T_k\left(\tfrac{x}{\varepsilon}\right)\, \partial_{x_k} v_0(x) \quad \text{and} \quad w_\varepsilon(x) = u_\varepsilon(x) - \overline{u}_\varepsilon(x) - \psi_\varepsilon(x), \quad x \in \overline{\Omega}_\varepsilon.$$

It is easy to verify that $supp\,(\psi_\varepsilon) \subset \mathcal{U}_{2\varepsilon} = \{x \in \overline{\Omega}_\varepsilon : \ \ \mathrm{dist}(x, \partial\Omega) \le 2\varepsilon\}$ and w_ε is a solution to the following problem

$$
\begin{cases}
-\mathcal{L}_\varepsilon(w_\varepsilon) = f_\varepsilon - f_0 - \sum_{m=1}^{2} q_m \left(g_0^{(m)}(x) - \kappa_m(v_0)\right) + \varepsilon\partial_{x_i}\left(F_i^\varepsilon(x)\right) + \mathcal{L}_\varepsilon(\psi_\varepsilon) + \\
+ \left(a_{ij}(\xi) + a_{ik}(\xi)\partial_{\xi_k}T_j(\xi) - |Q_0|^{-1}\hat{a}_{ij}\right)|_{\xi=\frac{x}{\varepsilon}} \ \partial^2_{x_ix_j}v_0 \qquad \text{in } \ \Omega_\varepsilon; \\
\sigma_\varepsilon(w_\varepsilon) = -\varepsilon\kappa_m(u_\varepsilon) + \varepsilon g_\varepsilon^{(m)}(x) - F_i^\varepsilon(x)\,\nu_i - \sigma_\varepsilon(\psi_\varepsilon) \qquad \text{on } \ \Xi_\varepsilon^{(m)} \quad (m = 1, 2); \\
w_\varepsilon = 0 \qquad \text{on } \ \Gamma_\varepsilon.
\end{cases}
$$

Multiplying the equation of this problem by w_ε, then integrating by parts and subtracting identities (9) for $\varphi_m = \kappa_m(v_0)\, w_\varepsilon, \ m = 1, 2$, we get

$$
\begin{aligned}
\varepsilon\sum_{m=1}^{2}\int_{\Xi_\varepsilon^{(m)}}\left(\kappa_m(u_\varepsilon) - \kappa_m(v_0)\right) w_\varepsilon\, ds_x + \int_{\Omega_\varepsilon} a_{ij}^\varepsilon\, \partial_{x_j}w_\varepsilon\, \partial_{x_i}w_\varepsilon\, dx = \\
= \int_{\Omega_\varepsilon}(f_\varepsilon - f_0)\varphi\, dx + \sum_{m=1}^{2}\Big(\varepsilon\int_{\Xi_\varepsilon^{(m)}} g_\varepsilon^{(m)}\, w_\varepsilon\, ds_x - q_m\int_{\Omega_\varepsilon} g_0^{(m)}\, w_\varepsilon\, dx\Big) - \\
- \varepsilon\sum_{m=1}^{2}\int_{\Omega_\varepsilon} a_{ij}^\varepsilon(x)\, \partial_{\xi_j}\psi_0|_{\xi=\frac{x}{\varepsilon}}\partial_{x_i}\left(\kappa_m(v_0)\, w_\varepsilon\right)\, dx + \\
+ \int_{\Omega_\varepsilon}\left(a_{ij}(\xi) + a_{ik}(\xi)\partial_{\xi_k}T_j(\xi) - |Q_0|^{-1}\widehat{a}_{ij}\right)|_{\xi=\frac{x}{\varepsilon}} \ \partial^2_{x_ix_j}v_0\, w_\varepsilon\, dx + \\
+ \varepsilon\int_{\Omega_\varepsilon} F_i^\varepsilon\partial_{x_i}w_\varepsilon\, dx - \int_{\Omega_\varepsilon} a_{ij}^\varepsilon\, \partial_{x_j}\psi_\varepsilon\, \partial_{x_i}w_\varepsilon\, dx. \qquad (51)
\end{aligned}
$$

Due to (1), (3) and (14) the left-hand side of (51) is estimated by the following way

$$
\begin{aligned}
\int_{\Omega_\varepsilon} a_{ij}^\varepsilon\, \partial_{x_j}w_\varepsilon\, \partial_{x_i}w_\varepsilon\, dx + \varepsilon\sum_{m=1}^{2}\int_{\Xi_\varepsilon^{(m)}}\left(\kappa_m(u_\varepsilon) - \kappa_m(v_0)\right) w_\varepsilon\, ds_x \ge \\
\ge c_1\Big(\int_{\Omega_\varepsilon}|\nabla w_\varepsilon|^2\, dx + \varepsilon\int_{\Xi_\varepsilon} w_\varepsilon^2\, ds_x\Big) - c_2\varepsilon\int_{\Xi_\varepsilon}\left|\left(\varepsilon T_k\, \partial_{x_k}v_0 + \psi_\varepsilon\right)w_\varepsilon\right|\, ds_x \ge \\
\ge c_3\|w_\varepsilon\|^2_{H^1(\Omega_\varepsilon)} - c_2\varepsilon\int_{\Xi_\varepsilon}\left|\left(\varepsilon T_k\, \partial_{x_k}v_0 + \psi_\varepsilon\right)w_\varepsilon\right|\, ds_x. \qquad (52)
\end{aligned}
$$

Now estimate the summands in the right-hand side of (51). Evidently, $|\int_{\Omega_\varepsilon}(f_\varepsilon - f_0)\, w_\varepsilon\, dx| \le \|f_\varepsilon - f_0\|_{L^2(\Omega_\varepsilon)}\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)}$. With the help of (9), (2) and (10) we bound the second and third terms:

$$
\begin{aligned}
\Big|\sum_{m=1}^{2}\Big(\varepsilon\int_{\Xi_\varepsilon^{(m)}} g_\varepsilon^{(m)}\, w_\varepsilon\, ds_x - q_m\int_{\Omega_\varepsilon} g_0^{(m)}\, w_\varepsilon\, dx\Big)\Big| = \sum_{m=1}^{2}\Big(\varepsilon\Big|\int_{\Omega_\varepsilon} a_{ij}^\varepsilon\, \partial_{\xi_j}\psi_0(\xi)\big|_{\xi=\frac{x}{\varepsilon}}\, \partial_{x_i}(g_\varepsilon^{(m)}w_\varepsilon)\, dx\Big| + \\
+ q_m\Big|\int_{\Omega_\varepsilon} g_\varepsilon^{(m)}w_\varepsilon\, dx - \int_{\Omega_\varepsilon} g_0^{(m)}w_\varepsilon\, dx\Big|\Big) = c_1\varepsilon\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)} + c_2\sum_{m=1}^{2}\|g_\varepsilon^{(m)} - g_0^{(m)}\|_{L^2(\Omega_\varepsilon)}\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)};
\end{aligned}
$$

$$
\varepsilon\sum_{m=1}^{2}\Big|\int_{\Omega_\varepsilon} a_{ij}^\varepsilon(x)\, \partial_{\xi_j}\psi_0\big|_{\xi=\frac{x}{\varepsilon}}\, \partial_{x_i}\left(\kappa_m(v_0)w_\varepsilon\right)\, dx\Big| \le
$$

$$\le \varepsilon c_3 \int_{\Omega_\varepsilon} |\nabla v_0|\,|w_\varepsilon|\,dx + \varepsilon c_4 \sum_{m=1}^{2} \int_{\Omega_\varepsilon} |\kappa_m(v_0)|\,|\nabla w_\varepsilon|\,dx \le \varepsilon c_5\,\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)}.$$

Thank to (25) and the fact that the vector-functions (35) are weak solenoidal in $\square$, it follows from Lemma 16.4 ([3]) that

$$\left|\int_{\Omega_\varepsilon} \Big(a_{ij}(\xi) + a_{ik}(\xi)\partial_{\xi_k}T_j(\xi) - \tfrac{1}{|Q_0|}\,\widehat{a}_{ij}\Big)\,\big|_{\xi=\frac{x}{\varepsilon}}\ \partial^2_{x_i x_j} v_0\, w_\varepsilon\,dx\right| \le \varepsilon c_6\,\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)}.$$

It is easy to see that $\varepsilon|\int_{\Omega_\varepsilon} F^\varepsilon_i \partial_{x_i} w_\varepsilon\,dx| \le \varepsilon c_6\,\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)}$. The last summand in (51) is estimated with the help of Lemma 1.5 ([5]) and (50):

$$\varepsilon\left|\int_{\Omega_\varepsilon} a^\varepsilon_{ij}\,\partial_{x_j}\psi_\varepsilon\,\partial_{x_i} w_\varepsilon\,dx\right| = \varepsilon\left|\int_{\mathcal{U}_{2\varepsilon}} a^\varepsilon_{ij}\,\partial_{x_j}\psi_\varepsilon\,\partial_{x_i} w_\varepsilon\,dx\right| \le$$

$$\le c_7 \int_{\mathcal{U}_{2\varepsilon}} |\nabla v_0|\,|\nabla w_\varepsilon|\,dx + \varepsilon^2 c_8 \int_{\mathcal{U}_{2\varepsilon}} |D^2 v_0|\,|\nabla w_\varepsilon|\,dx \le$$

$$\le c_7\|v_0\|_{H^1(\mathcal{U}_{2\varepsilon})}\|w_\varepsilon\|_{H^1(\mathcal{U}_{2\varepsilon})} + \varepsilon^2 c_8\|v_0\|_{H^2(\Omega)}\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)} \le c_9\varepsilon^{\frac{1}{2}}\|v_0\|_{H^2(\Omega)}\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)}. \tag{53}$$

It is remain to bound the last term in (52). Thanks to (9) and (10) we have

$$\varepsilon^2 \int_{\Xi_\varepsilon} |T_k\,\partial_{x_k} v_0\, w_\varepsilon|\,ds_x \le 2\varepsilon^2 \int_{\Omega_\varepsilon} \Big|a^\varepsilon_{ij}\,\partial_{\xi_j}\psi_0(\xi)\big|_{\xi=\frac{x}{\varepsilon}}\,\partial_{x_i}\big(T_k\,\partial_{x_k} v_0\, w_\varepsilon\big)\Big|\,dx +$$

$$+ c_{10}\,\varepsilon \int_{\Omega_\varepsilon} \Big|T_k\,\partial_{x_k} v_0\, w_\varepsilon\Big|\,dx \le c_{11}\varepsilon\|v_0\|_{H^2(\Omega)}\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)}. \tag{54}$$

With the same arguments as in (53) and (54) we have

$$\varepsilon \int_{\Xi_\varepsilon} |\psi_\varepsilon\, w_\varepsilon|\,ds_x \le c_{12}\varepsilon^{\frac{1}{2}}\|v_0\|_{H^2(\Omega)}\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)}.$$

Finally, we conclude from (51), (52) and estimates obtained above that

$$\|w_\varepsilon\|_{H^1(\Omega_\varepsilon)} \le C_1\Big(\varepsilon^{\frac{1}{2}} + \|f_\varepsilon - f_0\|_{L^2(\Omega_\varepsilon)} + \sum_{m=1}^{2} \|g^{(m)}_\varepsilon - g^{(m)}_0\|_{L^2(\Omega_\varepsilon)}\Big). \tag{55}$$

Since $\|\psi_\varepsilon\|_{H^1(\Omega_\varepsilon)}$ is bounded above by $C_2\varepsilon^{\frac{1}{2}}$, we have from (55) that

$$\|u_\varepsilon - \overline{u}_\varepsilon\|_{H^1(\Omega_\varepsilon)} \le C_3\Big(\varepsilon^{\frac{1}{2}} + \|f_\varepsilon - f_0\|_{L^2(\Omega_\varepsilon)} + \sum_{m=1}^{2} \|g^{(m)}_\varepsilon - g^{(m)}_0\|_{L^2(\Omega_\varepsilon)}\Big), \tag{56}$$

where the constant C_3 is independent of ε.

Thus, we have proved the following result.

Theorem 3. *Between the solution u_ε to problem (4) and the approximation function (46) the estimate (56) holds.*

With the help of the approximation function (46) and estimate (56) we can obtain an estimate for the energy integrals.

Corollary 1. *The following estimate*

$$\big|E_\varepsilon(u_\varepsilon)-E_0(v_0)\big|\le C\Big(\varepsilon^{\frac12}+\|f_\varepsilon-f_0\|_{L^2(\Omega_\varepsilon)}+\sum_{m=1}^{2}\|g_\varepsilon^{(m)}-g_0^{(m)}\|_{L^2(\Omega_\varepsilon)}\Big). \tag{57}$$

is satisfied, where the energy integrals $E_\varepsilon(u_\varepsilon)$ and $E_0(v_0)$ are defined in (29).

Proof. By virtue of (56) we have

$$\partial_{x_i}u_\varepsilon=\partial_{x_i}v_0+\partial_{\xi_i}T_k(\xi)|_{\xi=\frac{x}{\varepsilon}}\,\partial_{x_k}v_0+r_i^\varepsilon(x),$$

where

$$\|r_i^\varepsilon\|_{L^2(\Omega_\varepsilon)}\le C_1\Big(\varepsilon^{\frac12}+\|f_\varepsilon-f_0\|_{L^2(\Omega_\varepsilon)}+\sum_{m=1}^{2}\|g_\varepsilon^{(m)}-g_0^{(m)}\|_{L^2(\Omega_\varepsilon)}\Big).$$

Then

$$\int\limits_{\Omega_\varepsilon}a_{ij}^\varepsilon\partial_{x_j}u_\varepsilon\,\partial_{x_i}u_\varepsilon\,dx=\int\limits_{\Omega_\varepsilon}a_{ij}^\varepsilon\big(\partial_{x_i}v_0+\partial_{\xi_i}T_k(\xi)\big|_{\xi=\frac{x}{\varepsilon}}\,\partial_{x_k}v_0\big)\big(\partial_{x_j}v_0+\partial_{\xi_j}T_l(\xi)\big|_{\xi=\frac{x}{\varepsilon}}\,\partial_{x_l}v_0\big)dx+p_\varepsilon, \tag{58}$$

where

$$p_\varepsilon=2\int_{\Omega_\varepsilon}a_{ij}^\varepsilon\left(\partial_{x_i}v_0+\partial_{\xi_i}T_k(\xi)|_{\xi=\frac{x}{\varepsilon}}\,\partial_{x_k}v_0\right)r_\varepsilon^j\,dx+\int_{\Omega_\varepsilon}a_{ij}^\varepsilon r_\varepsilon^i r_\varepsilon^j\,dx.$$

Taking into account the boundedness of a_{ij}^ε and $\partial_{\xi_j}T_l(\xi)$ and estimate (56), we get

$$\begin{aligned}|p_\varepsilon|&\le c_1\left(\|v_0\|_{H^1(\Omega)}\left(\int_{\Omega_\varepsilon}r_\varepsilon^i r_\varepsilon^i\,dx\right)^{\frac12}+\int_{\Omega_\varepsilon}r_\varepsilon^i r_\varepsilon^i\,dx\right)\le\\&\le c_2\left(\varepsilon^{\frac12}+\|f_\varepsilon-f_0\|_{L^2(\Omega_\varepsilon)}+\sum_{m=1}^{2}\|g_\varepsilon^{(m)}-g_0^{(m)}\|_{L^2(\Omega_\varepsilon)}\right).\end{aligned} \tag{59}$$

Due to (2) we can regard here that $\|f_\varepsilon-f_0\|^2_{L^2(\Omega_\varepsilon)}\le\|f_\varepsilon-f_0\|_{L^2(\Omega_\varepsilon)}$, similar for other summands.

Let us introduce the following functions

$$H_{kl}(\xi)\equiv a_{ij}(\xi)\,\partial_{\xi_i}\left(T_k(\xi)+\xi_k\right)\partial_{\xi_j}\left(T_l(\xi)+\xi_l\right)\;-\;\tfrac{1}{|Q_0|}\,\widehat{a}_{kl},\quad k,l=1,\dots,n.$$

After extending the functions $a_{ij},\ T_k,\ \partial_{\xi_i}T_k,\ k=1,\dots,n$, by zero to $\square\backslash Q_0$, the functions $H_{kl},\ k,l=1,\dots,n$, will be 1-periodic with zero average over $\square$.

By the same way as we rewrote a summand in (41) (see (42)) and using the functions $H_{kl},\ k,l=1,\dots,n$, and (58), we obtain

$$\begin{aligned}\int_{\Omega_\varepsilon}a_{ij}^\varepsilon\partial_{x_j}u_\varepsilon\,\partial_{x_i}u_\varepsilon\,dx-\int_{\Omega}\widehat{a}_{ij}\partial_{x_j}v_0\,\partial_{x_i}v_0\,dx=\int_{\Omega}H_{kl}(\xi)|_{\xi=\frac{x}{\varepsilon}}\,\partial_{x_k}v_0\,\partial_{x_l}v_0\,dx+\\+\int_{\Omega}\big(\tfrac{1}{|Q_0|}\chi_{Q_0}(\tfrac{x}{\varepsilon})-1\big)\widehat{a}_{ij}\partial_{x_j}v_0\,\partial_{x_i}v_0\,dx+p_\varepsilon=:I_1+I_1+p_\varepsilon,\end{aligned} \tag{60}$$

where χ_{Q_0} is the characteristic function defined in (21). The summand I_1 can be estimated by the same way as in the proof of Theorem 1.3 ([5, Ch. 2]). As a result, we have $|I_1|\le c_1\varepsilon\|v_0\|^2_{H^2(\Omega)}$.

To estimate I_2 we note that $\int_\square\big(\frac{1}{|Q_0|}\chi_{Q_0}(\xi)-1\big)d\xi=0$. Therefore, with the help of Lemma 1.1 ([3]) we get $|I_2|\le c_2\varepsilon\|v_0\|^2_{H^2(\Omega)}$.

Summarizing (59) and estimates for I_1 and I_2, from (60) we deduce as follows

$$\Big|\int_{\Omega_\varepsilon} a^\varepsilon_{ij}\partial_{x_j}u_\varepsilon\,\partial_{x_i}u_\varepsilon\,dx-\int_\Omega \widehat{a}_{ij}\partial_{x_j}v_0\,\partial_{x_i}v_0\,dx\Big|\le$$

$$\le C_1\Big(\varepsilon^{\frac12}+\varepsilon\|f_0\|^2_{L^2(\Omega)}+\|f_\varepsilon-f_0\|_{L^2(\Omega_\varepsilon)}+\sum_{m=1}^{2}\|g^{(m)}_\varepsilon-g^{(m)}_0\|_{L^2(\Omega_\varepsilon)}\Big).\quad(61)$$

Now consider the difference

$$I_3:=\varepsilon\sum_{m=1}^{2}\int_{\Xi^{(m)}_\varepsilon}k_m(u_\varepsilon)u_\varepsilon\,ds_x-\sum_{m=1}^{2}|S^m|\int_\Omega k_m(v_0)v_0\,dx.$$

With the help of integral identities (9) we re-write it in the form

$$I_3=\varepsilon\sum_{m=1}^{2}\int_{\Omega_\varepsilon}a^\varepsilon_{ij}\,\partial_{\xi_j}\psi^{(m)}_0(\xi)\Big|_{\xi=\frac{x}{\varepsilon}}\partial_{x_i}(k_m(u_\varepsilon)u_\varepsilon)\,dx+$$

$$+\sum_{m=1}^{2}q_m\int_{\Omega_\varepsilon}k_m(u_\varepsilon)u_\varepsilon\,dx-\sum_{m=1}^{2}|S^m|\int_\Omega k_m(v_0)v_0\,dx.$$

Due (3), (10) and (17) the first term is not grater then εc_1. Since

$$\Big|\int_{\Omega_\varepsilon}k_m(u_\varepsilon)u_\varepsilon\,dx-\int_{\Omega_\varepsilon}k_m(v_0)v_0\,dx\Big|\le c_2\|u_\varepsilon-v_0\|_{L^2(\Omega_\varepsilon)},$$

it remains to estimate the following difference

$$\sum_{m=1}^{2}\Big|q_m\int_{\Omega_\varepsilon}k_m(v_0)v_0\,dx-|S^m|\int_\Omega k_m(v_0)v_0\,dx\Big|=\sum_{m=1}^{2}\Big|\int_\Omega\big(q_m\chi_{Q_0}(\tfrac{x}{\varepsilon})-|S^m|\big)k_m(v_0)v_0\,dx\Big|.$$

Thanks to the equality $\int_\square\big(q_m\chi_{Q_0}(\xi)-|S^m|\big)d\xi=0$ ($q_m=|S^m|/|Q_0|$) and Lemma 1.1 ([3]), this difference is bounded by $c_3\varepsilon\|v_0\|_{H^1(\Omega)}$. Thus, $|I_3|\le c_4\varepsilon+c_2\|u_\varepsilon-v_0\|_{L^2(\Omega_\varepsilon)}$.

Finally, taking into account the previous estimate, (61), (56) and noting that $E_\varepsilon(u_\varepsilon)-E_0(v_0)=I_1+I_2+I_3+p_\varepsilon$, we arrive to (57). □

References

[1] *Cioranescu D., Saint Jean Paulin J.* Homogenization of reticulated structures. Appl. Math. Sci. – New York: Springer-Verlag, 1999. – Vol. 136 – 346 p.

[2] *Cioranescu D., Donato P., Zaki R.* Asymptotic behavior of elliptic problems in perforated domains with nonlinear boundary conditions // Asymptotic Analysis. – 2007. – **53**, no. 4. – P. 209–235.

[3] *Chechkin G.A., Piatnitskii A.L., Shamaev A.S.* Homogenization: Methods and Applications. Translations of Mathematical Monographs. – Providence: AMS, 2007. – V. 234. – 234 p. (Translated from: Homogenization: Methods and Applications. – Novosibirsk: Tamara Rozhkovskaya Press, 2007. – 264 p.)

[4] *Marchenko V. A., Khruslov E. Ya.,* Homogenized models of micro-inhomogeneous medius. – Kyiv: Naukova Dumka, 2005. – 549 p. (in Russian).

[5] *Oleinik O.A., Yosifian G.A., Shamaev A.S.* Mathematical Problems in Elasticity and Homogenization. – Amsterdam: North-Holland, 1992. – 309 p. (Russian edition: Moscow: Izd-stvo MGU, 1990. – 310 p.)

[6] *Skrypnik I.V.* Methods for investigation of nonlinear elliptic boundary-value problems. – Moscow: Nauka, 1990. – 442 p. (in Russian).

[7] *Zhikov V.V., Kozlov S.M., Oleinik O.A.* Homogenization of differential operators and integral functionals. – Berlin: Springer-Verlag, 1994. – 570 p.

[8] *Zhikov V.V.* On the homogenization of nonlinear variational problems in perforated domains // Russian Journal of Mathematical Physics. – 1994. – **12**, no. 3. – P. 393–408.

[9] *Zhikov V.V.* Connectedness and homogenization. Examples of fractal conductivity // Mat. Sb. – 1996. – **187**, no. 8. – P. 3–40. in English: Sb. Math. – 1996. – **187**, no. 8. – P. 109–147.

[10] *Zhikov V.V., Rychago M.Ye.* Homogenization of nonlinear elliptic equations of the second order in perforated domains // Izv. Ross. Acad. Nauk, Ser. Mat. – 1997. – **61**, no. 1. – P. 69–89.

[11] *Khruslov E. Ya.* Asymptotic behaviour of solutions of the second order boundary value problem in the case of the refinement of the domain boundary // Mat. Sb. – 1978. – **106**, no. 4. – P. 604–621. (in Russian).

[12] *Cioranescu D., Saint Jean Paulin J.* Homogenization in open sets with holes // J. Math. Anal. Appl. – 1979. – **71**. – P. 590–607.

[13] *Mel'nyk T.A.* Homogenization of a boundary-value problem with a nonlinear boundary condition in a thick junction of type 3:2:1 // Mathematical Models and Methods in Applied Sciences. – 2008. – **31**, no. 9. – P. 1005–1027. Published online: `http://dx.doi.org/10.1002/mma.951`

[14] *Evans L.C.* Partial Differential Equations. Graduate Studies in Mathematics. – Providence: AMS, 1998. – Vol.19. – 662 p.